




ELSEVIER

www.sciencedirect.com
www.rbmonline.com



ARTICLE

Impact of whole systems traditional Chinese medicine on in-vitro fertilization outcomes




Lee E Hullender Rubin ^{a,b,c,*}, Michael S Opsahl ^d, Klaus E Wiemer ^d,
Scott D Mist ^b, Aaron B Caughey ^b

^a Oregon College of Oriental Medicine, 75 NW Couch St, Portland, OR, 97210, USA; ^b Oregon Health and Science University, Portland, OR, USA; ^c Portland Acupuncture Studio, Portland, OR, USA; ^d Poma Fertility, Northwest Center for Reproductive Sciences, Kirkland, WA, USA

* Corresponding author. E-mail address: lrubin@ocom.edu (LE Hullender Rubin).



Lee Hullender Rubin, DAOM, MS, LAc, FABORM, is a clinician and researcher specializing in reproductive medicine, women's health and vulvar pain. She graduated with her master's degree from Bastyr University in 2001, and her doctorate from the Oregon College of Oriental Medicine (OCOM) in 2009. Dr Hullender Rubin was OCOM's first postdoctoral research fellow funded by a National Institutes of Health educational grant. She teaches the doctoral infertility module and is Adjunct Research Faculty at OCOM, and Visiting Research Faculty at Oregon Health and Science University. She currently practices at her clinic, the Portland Acupuncture Studio, in Portland, Oregon.

Abstract Patients undergoing IVF may receive either acupuncture or whole-systems traditional Chinese medicine (WS-TCM) as an adjuvant IVF treatment. WS-TCM is a complex intervention that can include acupuncture, Chinese herbal medicine, dietary, lifestyle recommendations. In this retrospective cohort study, 1231 IVF patient records were reviewed to assess the effect of adjuvant WS-TCM on IVF outcomes compared among three groups: IVF with no additional treatment; IVF and elective acupuncture on day of embryo transfer; or IVF and elective WS-TCM. The primary outcome was live birth. Of 1069 non-donor cycles, WS-TCM was associated with greater odds of live birth compared with IVF alone (adjusted odds ratio [AOR] 2.09; 95% confidence interval [CI] 1.36 to 3.21), or embryo transfer with acupuncture only (AOR 1.62; 95% CI 1.04 to 2.52). Of 162 donor cycles, WS-TCM was associated with increased live births compared with all groups (odds Ratio [OR] 3.72; 95% CI 1.05 to 13.24, unadjusted) or embryo transfer with acupuncture only (OR 4.09; 95% CI: 1.02 to 16.38, unadjusted). Overall, IVF with adjuvant WS-TCM was associated with greater odds of live birth in donor and non-donor cycles. These results should be taken cautiously as more rigorous research is needed. 

© 2015 Reproductive Healthcare Ltd. Published by Elsevier Ltd. All rights reserved.

KEYWORDS: acupuncture, Chinese herbal medicine, embryo transfer, in-vitro fertilization, live births, traditional Chinese medicine

Introduction

Women undergoing IVF commonly pursue adjuvant complementary and alternative health approaches to support their cycles. Given that early studies have found that acupuncture significantly improves IVF pregnancy rates, patients increasingly choose adjuvant acupuncture to optimize their chances of pregnancy and live birth (De Lacey et al., 2009). Utilization rates are unclear. In a survey of 428 Northern Californian couples, 22% reported using acupuncture (Smith et al., 2010), but of 118 Boston-area women surveyed, 47% women reported using acupuncture and 17% used herbs during their assisted reproduction technique cycle (Domar et al., 2012). Another survey of 77 patients found that 92% reported using acupuncture to support their IVF cycle (Aelion et al., 2009). A patient may seek an acupuncturist's support by self-referral, or her IVF physician may refer her, but it is also possible that IVF physicians may not be aware of patient use of these additional therapies (Boivin and Schmidt, 2009).

Investigations into the effect of acupuncture on IVF outcomes largely focus on a limited number of standardized treatments, usually two, on or around embryo transfer compared with no acupuncture, sham acupuncture or a placebo needle. Initial studies suggested that acupuncture on or around embryo transfer improved IVF outcomes (Dieterle et al., 2006; Paulus et al., 2002; Smith et al., 2006; Westergaard et al., 2006), and, in a meta-analysis of seven trials, acupuncture increased the odds of clinical pregnancy by 65% (Manheimer et al., 2008). As more trials were completed, the equivocal effects were found when this narrow dose of only two or three acupuncture sessions were compared to controls. In a review of 14 trials, two to three acupuncture treatments administered around embryo transfer did not improve clinical pregnancy rates compared with controls (Cheong et al., 2013b). Another review of 16 trials confirmed these findings, but the covariate of baseline pregnancy rate was found to be a significant mediator of acupuncture's effect (Manheimer et al., 2013). Additionally, a clear need exists for an adequate acupuncture control, as penetrating or non-penetrating 'sham or placebo' acupuncture controls are likely to be impractical and may have physiological effects (Manheimer, 2011; Vickers et al., 2012).

With the advent of two standardized acupuncture sessions as a complete investigational intervention, acupuncture providers thereby question whether it is a sufficient dose (Craig et al., 2014; Shen et al., 2015). In clinical practice, acupuncture treatment is not standardized and instead considers the patient singularly; treatment is individualised. A course of treatment can range from six to 24 treatments depending on the complexity of the case. Furthermore, if an IVF patient seeks adjuvant acupuncture treatment in the period before the day of embryo transfer, it is possible that the patient, her partner, or both, will receive a complex intervention at the acupuncturist's office, such as whole-systems traditional Chinese medicine (WS-TCM). This is a multi-dimensional intervention that can include any combination of modalities classified under the system of traditional Chinese medicine. These include any combination of acupuncture (the insertion of sterile, filiform needles in the body) (Cochrane et al., 2014), moxibustion (the burning of processed herb, *artemesia argyi*, on or near the body) (Nedeljkovic et al., 2013), Chinese herbal medicine (Cao et al., 2013; Ried and

Stuart, 2011; Tan et al., 2012), Chinese medical massage with a tool (guasha) or without (tuina), Chinese medicine-based dietary recommendations, breathing exercises (Qi Gong), or movement exercises (Tai Chi) (Noll and Wilm, 2009). It may also include recommendations for vitamins, supplements, or both, depending on the training and licensure of the provider. Beyond a published case study (Hullender Rubin, 2010) and expert texts (Liang, 2003; Lyttleton, 2004; Noll and Wilm, 2009), the effectiveness of the multi-dimensional WS-TCM approach on IVF outcomes is unclear.

Given this background, we sought to compare the reproductive outcomes of women who elected WS-TCM treatment in addition to their usual IVF care, and compared them with those who received the usual IVF care alone and to those who received two standardized acupuncture treatments on the day of embryo transfer acupuncture only. Our main objective was to compare the three groups on the primary outcome of live birth.

Materials and methods

Study design

This was a retrospective cohort study of WS-TCM effects on IVF reproductive outcomes compared with two groups: those who received acupuncture treatments only on the day of embryo transfer (ACU); and those who received IVF usual care alone. The Oregon College of Oriental Medicine Institutional Review Board approved this study on 11 January 2011 (IRB reference number 09-028). Reproductive outcomes data were obtained from a single, private IVF centre, the Northwest Center for Reproductive Sciences (NCRS), Kirkland, Washington.

Patient criteria

Of 1509 patient charts, only NCRS patients who underwent IVF with fresh donor or non-donor embryos transferred between August 2005 and December 2010 were included, regardless of embryo quality. Additionally, women who underwent pre-implantation genetic screening testing or intracytoplasmic sperm injection cycles were also included. To further reduce selection bias, all biomedical diagnoses were included. Diminished ovarian reserve was diagnosed if FSH was 12 mIU/ml or more, antral follicle count total less than 10, and anti-Müllerian hormone less than 1.0 ng/ml, or any combination of these measures. Since the mean age of the WS-TCM was higher than the other groups, this most likely accounted for the increased incidence of diminished ovarian reserve diagnosis. All patients who underwent transfers with frozen embryos ($n = 251$) or embryos from frozen oocytes ($n = 27$, research cycles) were excluded.

All included patient data ($n = 1231$) were reviewed to identify patients who received ACU, and then reviewed a second time to identify and tabulate patients who received WS-TCM, as all WS-TCM patients received ACU as part of their treatment plans. The data were de-identified and assigned unique identifiers. The key was only available to the primary investigator. Visit data were then independently abstracted

from acupuncture clinic financial records and confirmed via scheduling calendars by two research assistants. A third party was consulted to resolve any conflicts.

Interventions

Usual care

Non-donor. IVF cycle management was individualized to each patient according to biomedical diagnosis. In general, patients were treated with either a gonadotrophin-releasing hormone (GnRH) agonist (Cramer et al., 1999) or antagonist (Felberbaum et al., 1995; Olivennes et al., 1998) IVF protocol. Oral contraceptive pills were taken on the first day of menses and continued from 1–3 weeks. Once ovarian suppression was confirmed, ovarian stimulation was initiated. Gonadotrophin medications were administered twice daily in variable doses based on age and antral follicle count, for up to 12 days, and included follitropin alpha (Gonal-f, EMD Serono, Geneva, Switzerland), follitropin beta injection (Follistim; Merck, New Jersey, US), injectable menotropins (Menopur, Ferring, Saint-Prex, Switzerland), or a combination of gonadotrophins. When three to four lead follicles measured a minimum of 17–18 mm on ultrasound, a patient was triggered with HCG (HCG; Novarel; Ferring, Siat-Prex, Switzerland). Within 36 h of trigger, transvaginal ultrasound-guided oocyte retrieval was carried out under conscious sedation. Once recovered from retrieval, patients were administered intramuscular progesterone in oil injections (Watson Laboratories, Parsippany, NJ, USA; West-Ward Pharmaceutical, Eatontown, NJ, USA; or Fresenius Kabi, Lake Zurich, IL, USA), 50 mg/ml twice daily until embryo transfer.

All recovered oocytes were fertilized within the embryology laboratory by either conventional insemination or intracytoplasmic sperm injection. Fertilization was visually confirmed by the presence of two pronuclei 17–22 h after fertilization. Embryos were cultured in 5% oxygen incubators and were evaluated every 24 h before embryo transfer. All embryos were individually cultured in 20–30 μ l droplets of Global Medium (IVF Online, Guildford, CT, USA), supplemented with 10% human serum albumin and covered with 5.5 ml of equilibrated oil. Embryo culture was carried out in 35 mm Nunc dishes (Thermo Scientific Fisher, San Rafael, CA, USA). For blastocyst culture, embryos were moved to new culture dishes on day 3 of development. Day of embryo transfer was based on the cause of patient infertility, number and quality of embryos, as well as patient age. Blastocyst culture and transfer was attempted whenever possible. When pre-implantation genetic diagnosis or screening was planned, blastomere biopsy was carried out on cleavage-stage embryos on day 3 of development, and embryo transfer occurred on day 5.

Day 3 embryo(s) or day 5 blastocyst(s) embryo transfer was undertaken with all patients with a full bladder using a Wallace catheter (Smiths Medical International, Kent, UK) with ultrasound guidance by one out of four physicians and according to SART guidelines (SART and ASRM, 2006; SART and ASRM, 2008; SART and ASRM, 2009). After transfer, Prometrium (progesterone, AbbVie Inc., Chicago, IL, USA), 200 mg suppositories were inserted vaginally twice daily. Serum HCG was collected to determine pregnancy 14 days after retrieval. If pregnant, progesterone supplementation continued to ges-

tational week 10. If not pregnant, progesterone was discontinued.

Donor. Egg donors underwent ovarian stimulation using similar protocols as non-donors. Donor egg recipients underwent an endometrial development cycle timed to correspond to the ovarian stimulation of the donor. Donor recipients began the oral contraceptive pill on the first day of their menses and continued from 1–3 weeks. Once suppression was confirmed, Vivelle-Dot patches (Novartis, Basel, Switzerland) and Estrace (Bayer-Schering, Berlin, Germany) were incrementally increased for up to 11–15 days depending on oestrogen levels and endometrial thickness. Once the endometrial thickness achieved 8 mm or thicker, and the donor eggs were retrieved, the donor recipient proceeded with the same progesterone protocol and embryo transfer as non-donor patients.

Acupuncture group

In addition to the IVF usual care, patients could elect to receive two standardized acupuncture treatments on the day of embryo transfer (ACU). The IVF nurse notified patients that acupuncture treatment was available to them. All ACU treatment protocols were based on previous published research (Craig et al., 2014; Paulus et al., 2002), and the treatment methods were previously described (Hullender Rubin et al., 2012a,b 2013c). Patients received two standardized acupuncture treatments on the same day as the embryo transfer.

Treatments were administered onsite at the IVF centre by one of nine state-licensed acupuncturists, certified in acupuncture by the National Certification Commission of Acupuncture and Oriental Medicine (NCCAOM). The acupuncturists' experience ranged from 1–9 years. All were trained to carry out the treatments in the same manner, and a treatment manual was provided. Refresher training to ensure continuity of methods occurred annually.

The first acupuncture session took place 1 h and 15 min before embryo transfer, and the second no more than 30 min after the embryo transfer. The Craig acupuncture protocol was used on both cycle types between 2005 and 2007 (Craig et al., 2014), and the following points were needled. Before embryo transfer: GV-20/Baihui; PC-6/Neiguan; CV-6/Qihai; ST-29/Guilai; SP-8/Diji; LR-3/Taichong; right ear (uterus, endocrine); left ear (Shenmen, Brain) After embryo transfer: LI-4/Hegu; SP-10/Xuehai; ST-36/Zusanli; SP-6/Sanyinjiao; KI-3/Taixi; right ear (Shenmen, Brain); and left ear (uterus, endocrine). Between 2008 and 2009, the modified Craig protocol was used in both cycle types, with the only modification being acupoint KI-3/Taixi was removed from the protocol (Hullender Rubin et al., 2012a,b). In 2010, the acupuncture protocol was adjusted according to cycle type owing to preliminary retrospective analysis suggesting fresh, non-donor cycles benefitted from the modified Craig (Hullender Rubin et al., 2012a), and donor cycles benefitted from the Craig protocol (Hullender Rubin et al., 2012b, 2013a). Patients could also elect to listen to soft music or a guided meditation during the acupuncture sessions. No other recommendations were provided.

Whole-systems traditional Chinese medicine group

In addition to the usual IVF care and acupuncture on the day of embryo transfer as described above, patients could elect

to pursue adjuvant whole-systems traditional Chinese Medicine (WS-TCM) treatment. Patients were either referred by the IVF doctor or self-referred. Flyers describing care with the acupuncturist were available in the IVF clinic lobby.

One of three Washington state-licensed acupuncturists with 3–4 years of master's level training in acupuncture and Chinese herbal medicine provided the WS-TCM treatment. All three acupuncturists were nationally certified by the NCCAOM in acupuncture, one (LHR) was NCCAOM certified in herbs, and one was certified in herbs by the California state licensing exam. All pursued postgraduate continuing education in reproductive medicine and infertility, and one (LHR) was board certified in reproductive medicine by the American Board of Oriental Reproductive Medicine.

At the initial visit, the patient's biomedical chart was reviewed. The patient was assessed according to TCM theory, an accepted form of medicine (WHO, 2007). A detailed WS-TCM treatment plan was provided outlining the dose and frequency of acupuncture, electro-acupuncture, Chinese medicine-based dietary suggestions, lifestyle recommendations, and any recommended nutritional supplements.

Patient visits were typically once a week and increased in frequency to twice a week from the start of IVF suppression medications to oocyte retrieval. Needles sizes ranged from 0.18 mm to 0.25 mm in diameter and 30–40 mm in length for body points, and 0.16 mm x 15 mm for ears. Size and length were selected based on patient's tolerance, acupoint location and body size. Needles used were Vinco brand (China), Sierin brand (Japan), and DBC brand (Korea). Electroacupuncture stimulator used was ES-2 (Heliomed, China) between 2005 and 2007, and Electrostimulator 4c.Pro (Pantheon Research, Venice, CA, USA) was used between 2007 and 2010. Treatment could also include the following modalities: warming Far Infrared Mineral lamp (TDP CQ-27, FIRARD II, China), Chinese medical massage (tuina) and indirect moxibustion stick (Hoist Wuyan Jiutiao, Smokeless Moxa Stick, China). All customized Chinese herb formulas were dispensed in granular form and manufactured by KPC herbs (Taiwan). All herbal tablets were made of standardized Chinese herbal formulas and manufactured by Golden Flower Chinese Herbs (Taiwan). Recommended nutritional supplements included prenatal vitamins, essential fatty acids, antioxidants and probiotics.

Acupuncture and herbal treatment was based on TCM pattern diagnosis and phase of menstrual cycle. According to Chinese medicine, regulating the menstrual cycle is considered a central treatment approach to female reproduction (Cochrane et al., 2014). This approach was previously described in several case studies (Hullender, 2009; Hullender Rubin, 2010; Hullender Rubin and Marx, 2012; Hullender Rubin et al., 2013b), but in general, treatment was divided into four phases: menses (cycle days 1–4, follicular [cycle days 5–11], ovulation [cycle days 12–16], and luteal [cycle days 17 to onset of menses]). During menses, treatment was directed at regulating menses and reducing of dysmenorrhoea pain, if present (Smith et al., 2011). Follicular phase treatment was directed at regulating reproductive hormones (Chen, 1997; Napadow et al., 2008), and endometrial (Stener-Victorin et al., 1996) and follicular development (Rashidi et al., 2013). Ovulation phase treatment was directed at inducing ovulation (Jedel et al., 2011; Johansson and Stener-Victorin, 2013; Johansson et al., 2013; Stener-Victorin et al., 2012). Luteal

phase treatment focused on regulating hormones (Cochrane et al., 2014). Throughout the entire cycle, treatment had a sub-aim of reducing stress (De Lacey et al., 2009; Kovarova et al., 2010; Madaschi et al., 2010). During the IVF cycle, WS-TCM treatments were individualized according to patient needs and TCM pattern diagnosis, but aimed to support the goals of the IVF phase. Treatment could also include Chinese medicine-based dietary and lifestyle recommendations. During down-regulation and ovarian stimulation, treatment was aimed to improve ovarian and uterine blood flow (Magarelli et al., 2009; Stener-Victorin et al., 1996), improve oocyte quality (Rashidi et al., 2013) and reduce stress. Acupuncture was optional on the day of oocyte retrieval to manage pain (Gejervall et al., 2005; Humaidan et al., 2006; Stener-Victorin, 2005), or postoperative nausea, vomiting, or both (Cheong et al., 2013a), which was used by only six patients. One more visit was suggested 5–7 days after embryo transfer to reduce stress.

Outcome measures

The primary outcome measure was live birth after 24 weeks' gestation. Secondary outcome measures were biochemical pregnancies, spontaneous abortions, ectopic pregnancies, gestational age, singleton, twin and triplet pregnancy. Biochemical pregnancy was defined as a positive serum beta-HCG and pregnancy that failed to develop to the clinically identifiable stage on ultrasound evaluation. Spontaneous abortion is defined as the spontaneous loss of a pregnancy with an ultrasound-confirmed gestational sac before 20 gestational weeks.

Statistical analysis

Treatment groups were counted and differentiated by donor or non-donor cycle type. Demographics and descriptive statistics were computed by group in the following variables for non-donor cycles: age, FSH, primary infertility, cause of infertility, cycle number, method of fertilization, pre-implantation genetic screening, total antral follicle count, number of days stimulated, total gonadotrophin dose, peak oestradiol, endometrial thickness, number of mature oocytes obtained, number of embryos transferred, day of embryo transfer (day 3 or 5), ovarian hyperstimulation syndrome (OHSS) and gestational age. For donor cycles, the same covariates were compared with the exception of age, FSH, total antral follicle count, number of days stimulated, total gonadotrophin dose and OHSS.

Outcomes of live birth, biochemical pregnancy, spontaneous abortion, singleton, twins and triplet pregnancy were tabulated by group and cycle type. Differences in outcomes were calculated using logistic regression and adjusted by covariates associated with outcomes.

Data were analysed in the following ways: (i) comparability of demographics between group; (ii) differences in outcomes with chi-squared (when groups numbered 10 or greater) or Fisher's exact (when groups numbered less than 10). Means were compared via analysis of variance and proportions were compared by chi-squared; and (iii) in order to control for potential confounding, the magnitude of effects on pregnancy

outcomes was assessed using multivariable logistic regression analyses and presented as adjusted odds ratios (AOR) with 95% confidence intervals (CI). Results were considered statistically significantly different if $P < 0.05$. Stata. (StataCorp, 2011, USA) was used to carry out all tests.

Results

A total of 1231 fresh cycles took place, in which an embryo transfer occurred. Non-donor cycles numbered 1069 and donor cycles numbered 162 (Table 1). In the non-donor cycles, 580

patients were in the usual care group, 370 in the ACU group and 119 in the WS-TCM group. The mean (\pm SD) number of WS-TCM visits was 12.0 (\pm 12.4). The three groups were comparable on all reproductive variables at the start of the cycle, except diagnosis of diminished ovarian reserve ($P = 0.01$), diagnosis of cause of infertility as 'other' ($P = 0.03$); cycle number ($P = 0.004$) and total antral follicle count ($P = 0.0497$). Significant differences were found in cycle characteristics: number of mature oocytes obtained ($P = 0.02$); number of embryos transferred ($P = 0.003$) (Table 2).

The proportion of live births was significantly higher in the WS-TCM group (61.3%) compared with either the usual care

Table 1 Demographic characteristics.

<i>Non-donor cycles</i>	<i>WS-TCM</i>	<i>Usual care</i>	<i>ACU</i>	<i>P-value</i>
Group N	119	580	370	
Age ^{a,b}	35.0 \pm 4.0	34.3 \pm 4.3	34.9 \pm 4.3	NS
FSH ^{a,b}	10.9 \pm 3.4	10.7 \pm 3.1	11.2 \pm 3.9	NS
Total antral follicle count ^{a,b,f}	8.0 \pm 2.4	8.3 \pm 2.5	7.9 \pm 2.6	0.0497
Primary infertility ^{c,d}	48.0	51.4	52.7	NS
Cause of infertility ^{c,d}				
Male	40.0	36.6	35.4	NS
Diminished ovarian reserve ^f	31.1	20.9	27.3	0.01
Endometriosis	10.1	11.9	12.7	NS
Ovulatory dysfunction	17.7	20.7	21.6	NS
Tubal	15.1	15.9	13.0	NS
Uterine	6.7	5.9	5.2	NS
Other ^f	13.5	7.8	6.0	0.03
Unknown	8.4	14.5	14.3	NS
Multiple diagnoses	34.5	31.4	30.8	NS
Cycle number ^{c,d}				
First cycle ^f	57.4	72.6	70.0	0.004
Repeated cycle ^f	42.6	27.4	30.0	0.004
<i>Donor cycles</i>	<i>WS-TCM</i>	<i>Usual care</i>	<i>ACU</i>	<i>P-value</i>
Group N	21	104	37	
Primary infertility ^{c,d}	47.6	34.6	45.9	NS
Cause of infertility				
Male ^{c,e}	19.0	16.3	10.8	NS
Diminished ovarian reserve ^{c,d}	81.0	92.3	89.2	NS
Endometriosis ^{c,e}	0.0	16.3	10.8	NS
Ovulatory dysfunction ^{c,e}	9.5	1.0	2.7	NS
Tubal ^{c,e}	9.5	6.7	5.4	NS
Uterine ^{c,e}	9.5	7.6	5.4	NS
Other ^{c,e}	4.8	9.6	8.1	NS
Unknown ^{c,e}	0.0	0.0	0.0	NS
Multiple diagnoses ^{c,e,f}	28.6	42.3	24.3	0.02
Cycle number				
First cycle ^{c,e}	57.1	51.0	67.6	NS
Repeated cycle ^{c,e}	42.9	49.0	32.4	NS

^aData are mean \pm SD.

^bAnalysis of variance.

^cData are proportion.

^dChi-squared.

^eFisher's exact.

^fSignificant findings ($P < 0.05$).

ACU = women receiving acupuncture on day of embryo transfer; NS = not statistically significant; WS-TCM = Whole-systems traditional Chinese medicine.

Table 2 Cycle characteristics.

	WS-TCM	Usual care	ACU	P-value
Non-donor cycles				
Group, n	119	580	370	
Number of days stimulated ^{a,b}	9.8 ± 1.2	9.9 ± 1.2	9.9 ± 1.3	NS
Total gonadotrophin dose, IU ^{a,b}	3959.3 ± 1503.6	3721.6 ± 1515.6	3932.1 ± 1580.1	NS
Peak oestradiol, pg/ml ^{a,b}	3153.7 ± 1781.8	3211.0 ± 1644.0	3017.0 ± 1634.0	NS
Endometrial thickness, mm ^{a,b}	10.9 ± 2.4	11.2 ± 2.4	11.2 ± 2.2	NS
Number of mature oocytes obtained ^{a,b,f}	14.0 ± 8.7	15.7 ± 9.1	14.2 ± 8.0	0.02
OHSS ^{c,d}	8.4	9.5	7.1	NS
Fertilization method ^{c,d}				
Conventional	47.1	51.0	46.3	NS
ICSI	48.7	46.2	49.3	NS
Mixed	4.2	2.8	4.3	NS
Pre-implantation genetic screening ^{c,d}	13.5	9.3	6.5	NS
Number of embryos transferred ^{a,b,f}	2.9 ± 1.2	2.6 ± 1.1	2.8 ± 1.1	0.003
Day of embryo transfer ^{c,d}				
Day 3	64.7	59.7	65.7	NS
Day 5	35.3	40.3	34.3	NS
Mean number of acupuncture visits ^a	12.0 ± 12.4			
Donor cycles				
Group, N	21	104	37	
Peak oestradiol, pg/ml ^{a,b}	4303.9 ± 2130.5	4250.3 ± 2035.4	4428.4 ± 2660.4	NS
Endometrial thickness, mm ^{a,b}	9.7 ± 2.0	10.5 ± 2.1	10.0 ± 2.0	NS
Number of mature oocytes obtained from donor ^{a,b,f}	18.6 ± 8.0	21.7 ± 8.7	26.4 ± 11.2	0.005
Fertilization method				
Conventional ^{c,d}	57.1	60.6	67.6	NS
ICSI ^{c,e}	38.1	33.7	27.0	NS
Mixed ^{c,e}	4.8	5.8	5.4	NS
Pre-implantation genetic diagnosis Testing ^{c,e}	4.8	10.6	8.3	NS
Number of embryos transferred ^{a,b}	2.2 ± 0.7	2.2 ± 0.7	1.9 ± 0.6	NS
Day of embryo transfer				
Day 3 ^{c,e}	23.8	38.5	24.3	NS
Day 5 ^{c,d}	76.2	61.5	76.7	NS
Mean number of acupuncture visits ^a	10.6 ± 13.6			

^aData are mean ± SD.

^bAnalysis of variance.

^cData are proportions.

^dChi-squared.

^eFisher exact.

^fSignificant findings ($P < 0.05$).

ACU = day of embryo transfer acupuncture only; ICSI = intracytoplasmic injection; NS = not significant; OHSS = ovarian hyperstimulation syndrome; WS-TCM = whole-systems traditional Chinese medicine.

(48.2%) or ACU groups (50.8%, $P = 0.03$ (Table 3)). Fewer spontaneous abortions ($P = 0.02$) occurred in the WS-TCM group compared with the other groups. On the outcome of biochemical pregnancies, there was no difference between groups, but WS-TCM trended toward significantly fewer than the other groups ($P = 0.05$). No ectopic pregnancies occurred in the WS-TCM group, but this was not a significant difference than compared with the usual care or ACU groups. Mean gestational age, and the proportions of live birth on or after 37 gestational weeks, singleton, twin and triplet pregnancies were not different between groups.

The WS-TCM group was associated with more live births compared with both groups when adjusted for the covariates of previous IVF cycle, age, and gonadotrophin dosage (AOR

1.93; 95% CI 1.27 to 2.92) (Table 4). More live births were associated with WS-TCM compared with embryo transfer acupuncture only, adjusted for age and gonadotrophin dosage (AOR 1.62; 95% CI 1.04 to 2.52), and when compared with IVF alone, adjusted for previous IVF and gonadotrophin dosage (AOR 2.09; 95% CI 1.36 to 3.21). Live births were not significantly affected in patients who received acupuncture only on the day of embryo transfer compared with IVF usual care.

When compared with both groups, WS-TCM was associated with fewer biochemical pregnancies (odds ratio [OR] 0.27; 95% CI 0.08 to 0.86). Decreased biochemical pregnancies were also associated with WS-TCM compared with embryo transfer acupuncture alone unadjusted (OR 0.25; 95% CI 0.07 to

Table 3 Reproductive outcomes.

<i>Non-donor cycles</i>	<i>WS-TCM</i>	<i>Usual care</i>	<i>ACU</i>	<i>P-value</i>
Group, <i>n</i>	119	580	370	
Live birth ^{a,b,f}	61.3	48.2	50.8	0.03
Biochemical pregnancy ^{a,c,f}	2.5	8.4	9.5	NS
Spontaneous abortion ^{a,c,f}	5.8	10.7	7.3	0.02
Ectopic ^{a,c}	0	1.0	0.8	NS
Singleton ^{a,b}	60.3	63.2	57.0	NS
Twins ^{a,b}	38.4	33.2	40.3	NS
Triplets ^{a,b}	1.4	3.2	2.2	NS
Gestational age ^{d,e}	37.8 ± 2.7	37.8 ± 2.8	37.4 ± 3.3	NS
Live birth on or after 37.0 weeks ^{a,b}	64.4	72.9	67.6	NS
<i>Donor cycles</i>				
Group, <i>n</i>	21	104	37	
Live birth ^{c,d}	85.7	62.5	59.5	NS
Biochemical pregnancy ^{a,c}	4.8	8.7	8.1	NS
Spontaneous abortion ^{a,c}	4.8	11.5	16.2	NS
Ectopic ^{a,c}	0.0	1.0	0.0	NS
Singleton ^{a,b}	55.6	54.0	54.6	NS
Twins ^{a,c}	38.9	45.3	45.5	NS
Triplets ^{a,c}	4.8	1.6	0.0	NS
Gestational age ^{d,e}	37.7 ± 3.3	36.9 ± 3.1	36.8 ± 3.7	NS
Live birth on or after 37.0 weeks ^{a,b}	72.2	60.0	63.6	NS

^aData are proportions.

^bChi-squared.

^cFisher's exact.

^dData are mean ± SD.

^eAnalysis of variance.

^fSignificant findings ($P < 0.05$).

ACU = day of embryo transfer acupuncture only; NS = not significant; WS-TCM = whole-systems traditional Chinese medicine.

Table 4 Reproductive outcomes logistic regression analysis (odds ratio and 95% confidence interval).

<i>Outcome</i>	<i>Cycle type</i>	<i>WS-TCM versus all</i>	<i>WS-TCM versus usual care</i>	<i>WS-TCM versus ACU</i>	<i>ACU versus usual care</i>
Live birth	Non-donor	1.93, 1.27 to 2.92 ^{a,h}	2.09, 1.36 to 3.21 ^{b,h}	1.62, 1.04 to 2.52 ^{c,h}	1.24, 0.94 to 1.64 ^d
	Donor	3.72, 1.05 to 13.24 ^{e,h}	3.6, 1.00 to 13.01 ^e	4.09, 1.02 to 16.38 ^{e,h}	0.74, 0.35 to 1.58 ^e
Biochemical pregnancy	Non-donor	0.27, 0.08 to 0.86 ^{e,h}	0.28, 0.09 to 0.91 ^{f,h}	0.25, 0.07 to 0.82 ^{e,h}	1.30, 0.83 to 2.03 ^e
	Donor	0.54, 0.07 to 4.36 ^e	0.55, 0.06 to 4.74 ^g	0.57, 0.06 to 5.82 ^e	1.01, 0.26 to 3.90 ^e
Spontaneous abortion	Non-donor	0.60, 0.27 to 1.34 ^e	0.50, 0.22 to 1.13 ^e	0.79, 0.34 to 1.87 ^e	0.72, 0.45 to 1.14 ^e
	Donor	0.34, 0.04 to 2.70 ^e	0.38, 0.05 to 3.12 ^e	0.26, 0.03 to 2.31 ^e	1.67, 0.59 to 4.75 ^e

^aAdjusted odds ratio comparing WS-TCM with all groups with previous IVF, age and gonadotrophin dosage as covariates.

^bAdjusted odds ratio comparing WS-TCM with usual care, with previous IVF and gonadotrophin dosage as covariates.

^cAdjusted odds ratio comparing WS-TCM with ACU with age and gonadotrophin dosage as covariates.

^dAdjusted odds ratio comparing usual care with ACU with prior IVF, age, and gonadotrophin dosage as covariates.

^eUnadjusted odds ratio.

^fAdjusted odds ratio comparing WS-TCM with usual care, with FSH, and gonadotrophin dosage as covariates.

^gAdjusted odds ratio comparing WS-TCM with usual care, with number of embryos transferred as covariate.

^hSignificant outcomes.

ACU = day of embryo transfer acupuncture only; usual care, usual IVF care; WS-TCM = whole-systems traditional Chinese medicine.

0.82), or standard IVF when adjusted for FSH and gonadotrophin dosage, (AOR 0.28; 95% CI 0.09 to 0.91). No difference was observed between the ACU and usual care groups on this outcome.

No differences were detected between groups on the outcomes of spontaneous abortion, ectopic pregnancies, singleton, twin, triplet pregnancies or gestational age.

Of the donor cycles, 104 recipients usual care group received an embryo transfer, 37 in the ACU group and 21 in the WS-TCM group. The mean (\pm SD) number of WS-TCM visits was 10.6 (\pm 13.6). The three groups were comparable on the variables of incidence of primary infertility; cause of infertility due to male factors; diagnosis of diminished ovarian reserve; endometriosis; ovulatory dysfunction; tubal issues; uterine issues; unknown reasons; other diagnosis; male only factors; female only factors; male and female factors; cycle number; incidence of pre-implantation genetic screening; fertilization method; number of days stimulated; peak oestradiol; endometrial thickness; day 3 or day 5 embryo transfer (**Tables 1 and 2**). Groups were different for the following variables: multiple diagnoses ($P = 0.02$) and number of mature oocytes obtained ($P = 0.005$).

On the outcome of live births, 85.7% of the WS-TCM group, 62.5% of the usual care and 59.5% of the ACU group had a live birth (**Table 3**). No differences were observed between groups for biochemical pregnancies or spontaneous abortions. No ectopic pregnancies occurred in the WS-TCM group for this cycle type, but this was not significantly fewer than usual care or ACU groups. Gestational age, singleton, twin, and triplet pregnancies were also not different between groups.

Significantly greater odds of live birth was associated with adjuvant WS-TCM compared with all groups (OR 3.72; 95% CI 1.05 to 13.24), and when compared with the ACU group (OR 4.09; 95% CI 1.02 to 16.38). When WS-TCM was compared with usual care, an association of increased odds trended towards significance (OR 3.6; 95% CI 1.00 to 13.01, unadjusted). The small sample size and exceedingly broad confidence intervals, however, indicate the weaknesses of these associations. No difference was found between the ACU and usual care groups on this outcome.

Outcomes of biochemical pregnancy or spontaneous abortion was not significantly affected by WS-TCM, regardless of comparison group in the donor cycles.

Discussion

To the best of our knowledge, this is the first adequately powered study of the effect of adjuvant WS-TCM on IVF reproductive outcomes. This is meaningful for its public health impact on infertility treatment and management, preliminary evidence of the need for an optimized acupuncture dose and demonstration that WS-TCM, a multidisciplinary intervention that includes acupuncture, was associated with more live births than two standardized acupuncture treatments on the day of embryo transfer or usual IVF care alone. These findings suggest WS-TCM could make significant contributions to the public health issue of infertility ([Macaluso et al., 2010](#); [CDC, 2010](#); [CDC, 2014](#)), if provided in an effective dose. Although IVF live births increase with advances in patient management and laboratory procedures, success is not guaranteed. Couples often must undergo several IVF cycles engendering extensive emotional distress and economic costs ([Cousineau and Domar, 2007](#)).

Acupuncture on or around day of embryo transfer does significantly reduce stress in IVF patients in both randomized controlled trials (RCTs) ([Domar et al., 2009](#); [Isoyama et al., 2012](#); [Smith et al., 2006](#)) and observational studies ([Balk et al., 2010](#); [De Lacey et al., 2009](#)). Initial RCTs found that acupuncture

improved IVF pregnancy rates ([Dieterle et al., 2006](#); [Paulus et al., 2002](#); [Westergaard et al., 2006](#)). As more studies were conducted, the effect became less clear as subsequent studies failed to reproduce those outcomes, with one RCT finding live births were lower in the acupuncture group compared with control ([Craig et al., 2014](#)). [Craig et al. \(2014\)](#) suspected the location of treatment offsite from the IVF centre and study population of acupuncture naïve women may have contributed to this finding (2014). Further, two standardized acupuncture sessions administered on day of embryo transfer alone were an insufficient acupuncture dose to improve IVF birth outcomes in two of the most recent systematic reviews ([Cheong et al., 2013b](#); [Manheimer et al., 2013](#)).

We observed that, in addition to embryo transfer acupuncture, an average of 12 WS-TCM visits for non-donor cycles and 11 visits for donor cycles were associated with greater odds of live birth. The combined approach of WS-TCM treatment before starting IVF with the two treatments on the day of embryo transfer acupuncture may be an optimal intervention to improve IVF birth outcomes in addition to stress reduction. An increased acupuncture dose, ie. more acupuncture sessions, is consistent with the effectiveness reported in other studies. Uterine blood flow was improved with eight electro-acupuncture sessions ([Stener-Victorin et al., 1996](#)). Nine electro-acupuncture sessions plus embryo transfer day acupuncture were correlated with an increase in serum cortisol and prolactin and improved IVF outcomes in prospective clinical trial ([Magarelli et al., 2009](#)). Fourteen sessions of low-frequency electro-acupuncture significantly reduced androgens and improved menstrual frequency over physical exercise or no treatment in women with polycystic ovary syndrome in an RCT ([Jedel et al., 2011](#)). Ovulation was induced in another RCT with 24 visits over 3 months in women with polycystic ovary syndrome ([Johansson et al., 2013](#)) using the same intervention described by [Jedel et al. \(2011\)](#).

In our study, every patient undergoing WS-TCM received a multidisciplinary intervention, of which all patients received acupuncture, electro-acupuncture, or both, in addition to lifestyle and dietary recommendations. Prenatal vitamins, essential fatty acids, antioxidants, probiotics, and Chinese herbs were also commonly suggested. Because of the nature of individualized care, recommendations were tailored to the patient's need and varied considerably within the group. Further, some patients elected to only pursue acupuncture to support their cycles. This is representative of TCM treatment in the real world. An objective of whole-systems research is to evaluate the combined effects of a complex intervention ([Elder et al., 2006](#); [Ritenbaugh et al., 2003](#)). It is not intended to evaluate the single ingredients of a complex intervention, as it may over or underestimate the single ingredient's influence and cannot account for the confounding or mediating effects of the other component(s).

This retrospective study has three limitations. First, owing to the nature of a retrospective cohort study, there is limited internal validity. Study groups were not randomly allocated, so the groups differed on several variables. We attempted to minimize potential bias from these differences by adjusting for covariates associated with pregnancy outcomes through multivariable logistic regression analysis. We also acknowledge the difference in the number of embryos transferred between groups may influence our findings. During dataset analysis, however, this covariate did not signifi-

cantly impact outcomes. We therefore question the biologic plausibility that the difference contributed, in part, to our results. Further, women elected which treatment to pursue. Women who elected WS-TCM, ACU, or both, could differ in unknown ways that may affect outcomes.

Secondly, it was also possible that women in the usual care groups received WS-TCM or day of embryo transfer acupuncture at another site, or patients in the ACU group received WS-TCM elsewhere leading to misclassification bias. Complementary and alternative medicine use is common in IVF patients (Smith et al., 2010) and not always reported to their IVF practitioners (Boivin and Schmidt, 2009). Such bias, however, would have been towards the null suggesting that our estimates of effect would only underestimate the actual effect of WS-TCM.

Finally, embryo quality may have differed between groups. Although embryo quality is a known predictor of IVF success (Heitmann et al., 2013; Vernon et al., 2011), only the highest quality embryos available were selected for transfer. As more women in the WS-TCM group were diagnosed with diminished ovarian reserve, it is more likely that the WS-TCM group would have had lower quality embryos, which may mean the effect of WS-TCM is underestimated by our study.

In conclusion, despite these potential limitations, this study provides evidence that the addition of WS-TCM to non-donor IVF cycles may increase the odds of achieving a live birth over usual IVF care alone or two standardized treatments administered around embryo transfer. The effects of WS-TCM were less clear compared with usual donor IVF care alone but showed a non-significant trend towards increased odds of a live birth in this cycle type. This retrospective cohort study suggests that WS-TCM as an adjuvant IVF treatment may be associated with improved live birth rates. More rigorous, comparative effectiveness research is needed to substantiate these findings and determine an optimal number of WS-TCM visits compared with an adequate and validated time-attention control, such as a mind-body programme to confirm these findings.

Acknowledgements

Sincere thanks to the NCRS patients, doctors, embryology, and nursing staff. LHR formerly owned the acupuncture practice. Whole-systems traditional Chinese medicine was provided by primary investigator (LHR); Roxane Geller, MS, LAc; and Holly Fritz, LAc. Day of embryo transfer acupuncture was provided by the primary investigator (LHR), HF; Melissa Rosenberger MS, MAcOM, LAc; Janci Karp, ND, LAc, LMP; Heather Stadler, MAC, LAc; Nicole Hidaka, MS, LAc; Lindsay A. Staker, MS, LAc; and Elaina Greenberg, MS, LAc. Chart data abstraction completed by Patrick Allen, MAcOM, LAc, and Angela Humphrey, MAcOM, LAc.

References

- Aelion, A., Barbieri, E., Shastri, S., Grill, E., Chung, P., Rosenwaks, Z., 2009. The use of complementary medical therapies (CMT) in infertility patients. *Fertil. Steril.* 92, 533–534.
- Balk, J., Catov, J., Horn, B., Gececi, K., Wakim, A., 2010. The relationship between perceived stress, acupuncture, and pregnancy rates among IVF patients: a pilot study. *Complement. Ther. Clin. Pract.* 16, 154–157.
- Boivin, J., Schmidt, L., 2009. Use of complementary and alternative medicines associated with a 30% lower ongoing pregnancy/live birth rate during 12 months of fertility treatment. *Hum. Reprod.* 24, 1626–1631.
- Cao, H., Han, M., Ng, E.H., Wu, X., Flower, A., Lewith, G., Liu, J.P., 2013. Can Chinese herbal medicine improve outcomes of in vitro fertilization? A systematic review and meta-analysis of randomized controlled trials. *PLoS ONE* 8, e81650.
- CDC, 2010. Outline for a National Action Plan for the Prevention, Detection and Management of Infertility. Centers for Disease Control and Prevention, Atlanta, GA.
- CDC, 2014. National Public Health Action Plan for the Detection, Prevention, and Management of Infertility. Centers for Disease Control and Prevention, Atlanta, GA.
- Chen, B.Y., 1997. Acupuncture normalizes dysfunction of hypothalamic-pituitary-ovarian axis. *Acupunct. Electrother. Res.* 22, 97–108.
- Cheong, K.B., Zhang, J.P., Huang, Y., Zhang, Z.J., 2013a. The effectiveness of acupuncture in prevention and treatment of post-operative nausea and vomiting—a systematic review and meta-analysis. *PLoS ONE* 8, e82474.
- Cheong, Y.C., Dix, S., Hung Yu Ng, E., Ledger, W.L., Farquhar, C., 2013b. Acupuncture and assisted reproductive technology. *Cochrane Database. Syst. Rev.* (7), CD006920.
- Cochrane, S., Smith, C.A., Possamai-Inesedy, A., Bensoussan, A., 2014. Acupuncture and women's health: an overview of the role of acupuncture and its clinical management in women's reproductive health. *Int. J. Womens Health* 6, 313–325.
- Cousineau, T.M., Domar, A.D., 2007. Psychological impact of infertility. *Best Pract. Res. Clin. Obstet. Gynaecol.* 21, 293–308.
- Craig, L.B., Hullender Rubin, L., Peck, J.D., Anderson, M., Marshall, L.A., Soules, M.R., 2014. Acupuncture performed before and after embryo transfer: a randomized controlled trial. *J. Reprod. Med.* 59, 313–320.
- Cramer, D.W., Powers, D.R., Oskowitz, S.P., Liberman, R.F., Hornstein, M.D., McShane, P.M., Barbieri, R.L., 1999. Gonadotropin-releasing hormone agonist use in assisted reproduction cycles: the influence of long and short regimens on pregnancy rates. *Fertil. Steril.* 72, 83–89.
- De Lacey, S., Smith, C.A., Paterson, C., 2009. Building resilience: a preliminary exploration of women's perceptions of the use of acupuncture as an adjunct to In Vitro Fertilisation. *BMC Complement. Altern. Med.* 9, 50.
- Dieterle, S., Ying, G., Hatzmann, W., Neuer, A., 2006. Effect of acupuncture on the outcome of in vitro fertilization and intracytoplasmic sperm injection: a randomized, prospective, controlled clinical study. *Fertil. Steril.* 85, 1347–1351.
- Domar, A.D., Meshay, I., Kelliher, J., Alper, M., Powers, R.D., 2009. The impact of acupuncture on in vitro fertilization outcome. *Fertil. Steril.* 91, 723–726.
- Domar, A.D., Conboy, L., Denardo-Roney, J., Rooney, K.L., 2012. Lifestyle behaviors in women undergoing in vitro fertilization: a prospective study. *Fertil. Steril.* 97, 697–701, e1.
- Elder, C., Aickin, M., Bell, I.R., Fonnebo, V., Lewith, G.T., Ritenbaugh, C., Verhoef, M., 2006. Methodological challenges in whole systems research. *J. Altern. Complement. Med.* 12, 843–850.
- Felberbaum, R.E., Reissmann, T., Kupker, W., Bauer, O., Al Hasani, S., Diedrich, C., Diedrich, K., 1995. Preserved pituitary response under ovarian stimulation with HMG and GnRH antagonists (Cetrorelix) in women with tubal infertility. *Eur. J. Obstet. Gynecol. Reprod. Biol.* 61, 151–155.
- Gejervall, A.L., Stener-Victorin, E., Moler, A., Janson, P.O., Werner, C., Bergh, C., 2005. Electro-acupuncture versus conventional analgesia: a comparison of pain levels during oocyte aspiration and patients' experiences of well-being after surgery. *Hum. Reprod.* 20, 728–735.
- Heitmann, R.J., Hill, M.J., Richter, K.S., Decherney, A.H., Widra, E.A., 2013. The simplified SART embryo scoring system is highly

- correlated to implantation and live birth in single blastocyst transfers. *J. Assist. Reprod. Genet.* 30, 563–567.
- Hullender, L., 2009. Vaginismus successfully treated with Chinese medicine – a case study. *Am. Acupunct.* 49, 24–27.
- Hullender Rubin, L., 2010. Adjunctive role of Chinese herbs and acupuncture in an in vitro fertilization cycle – a case study. *Am. Acupunct.* 52, 16–19.
- Hullender Rubin, L.E., Marx, B.L., 2012. Diminished ovarian reserve, clomid and traditional Chinese medicine – a case study. *Med. Acupunct.* 24, 273–280.
- Hullender Rubin, L.E., Opsahl, M.S., Ackerman, D.L., 2012a. P02.15. Acupuncture improves in vitro fertilization live birth outcomes – a retrospective chart review [Abstract]. *BMC Complement. Altern. Med.* 12, P71.
- Hullender Rubin, L.E., Opsahl, M.S., Ackerman, D.L., 2012b. P02.163. Acupuncture and donor egg in vitro fertilization cycles – a retrospective chart review comparing two acupuncture protocols. [Abstract]. *BMC Complement. Altern. Med.* 12, P219.
- Hullender Rubin, L., Opsahl, M.S., Wiemer, K., Ackerman, D.L., 2013a. P-5. A retrospective chart review comparing two acupuncture protocols on the day of in vitro fertilization fresh embryo transfer: do the points matter? [abstract]. *J. Altern. Complement. Med.* 19, A1.
- Hullender Rubin, L.E., Cantor, D., Marx, B.L., 2013b. Recurrent pregnancy loss and traditional Chinese medicine – a case study. *Med. Acupunct.* 25, 232–237.
- Hullender Rubin, L.E., Opsahl, M.S., Taylor-Swanson, L., Ackerman, D.L., 2013c. Acupuncture and in vitro fertilization live births – a retrospective chart review. *J. Altern. Complement. Med.* 19, 637–643.
- Humaidan, P., Brock, K., Bungum, L., Stener-Victorin, E., 2006. Pain relief during oocyte retrieval – exploring the role of different frequencies of electro-acupuncture. *Reprod. Biomed. Online* 13, 120–125.
- Isoyama, D., Cordts, E.B., de Souza van Niewegen, A.M., de Almeida Pereira de Carvalho, W., Matsumura, S.T., Barbosa, C.P., 2012. Effect of acupuncture on symptoms of anxiety in women undergoing in vitro fertilisation: a prospective randomised controlled study. *Acupunct. Med.* 30, 85–88.
- Jedel, E., Labrie, F., Oden, A., Holm, G., Nilsson, L., Janson, P.O., Lind, A.K., Ohlsson, C., Stener-Victorin, E., 2011. Impact of electro-acupuncture and physical exercise on hyperandrogenism and oligo/amenorrhoea in women with polycystic ovary syndrome: a randomized controlled trial. *Am. J. Physiol. Endocrinol. Metab.* 300, E37–E45.
- Johansson, J., Stener-Victorin, E., 2013. Polycystic ovary syndrome: effect and mechanisms of acupuncture for ovulation induction. *Evid. Based Complement. Alternat. Med.* 2013, 762615.
- Johansson, J., Redman, L., Veldhuis, P.P., Sazonova, A., Labrie, F., Holm, G., Johansson, G., Stener-Victorin, E., 2013. Acupuncture for ovulation induction in polycystic ovary syndrome: a randomized controlled trial. *Am. J. Physiol. Endocrinol. Metab.* 304, E934–E943.
- Kovarova, P., Smith, C.A., Turnbull, D.A., 2010. An exploratory study of the effect of acupuncture on self-efficacy for women seeking fertility support. *Explore (NY)* 6, 330–334.
- Liang, L., 2003. *Acupuncture and IVF*. Blue Poppy Press, Boulder, CO.
- Lyttleton, J., 2004. *Treatment of Infertility with Chinese Medicine*. Churchill Livingstone, Edinburgh.
- Macaluso, M., Wright-Schnapp, T.J., Chandra, A., Johnson, R., Satterwhite, C.L., Pulver, A., Berman, S.M., Wang, R.Y., Farr, S.L., Pollack, L.A., 2010. A public health focus on infertility prevention, detection, and management. *Fertil. Steril.* 93, e1–e10.
- Madaschi, C., Braga, D.P.A.F., Figueira, R.D.C.S., Iaconelli, A., Jr., Borges, E., Jr., 2010. Effect of acupuncture on assisted reproduction treatment outcomes. *Acupunct. Med.* 28, 180–184.
- Magarelli, P.C., Cridennda, D.K., Cohen, M., 2009. Changes in serum cortisol and prolactin associated with acupuncture during controlled ovarian hyperstimulation in women undergoing in vitro fertilization-embryo transfer treatment. *Fertil. Steril.* 92, 1870–1879.
- Manheimer, E., 2011. Selecting a control for in vitro fertilization and acupuncture randomized controlled trials (RCTs): how sham controls may unnecessarily complicate the RCT evidence base. *Fertil. Steril.* 95, 2456–2461.
- Manheimer, E., Zhang, G., Udoff, L., Haramati, A., Langenberg, P., Berman, B.M., Bouter, L.M., 2008. Effects of acupuncture on rates of pregnancy and live birth among women undergoing in vitro fertilisation: systematic review and meta-analysis. *BMJ* 336, 545–549.
- Manheimer, E., Van Der Windt, D., Cheng, K., Stafford, K., Liu, J., Tierney, J., Lao, L., Berman, B.M., Langenberg, P., Bouter, L.M., 2013. The effects of acupuncture on rates of clinical pregnancy among women undergoing in vitro fertilization: a systematic review and meta-analysis. *Hum. Reprod. Update* 19, 696–713.
- Napadow, V., Ahn, A., Longhurst, J., Lao, L., Stener-Victorin, E., Harris, R., Langevin, H.M., 2008. The status and future of acupuncture mechanisms research. *J. Altern. Complement. Med.* 14, 861–869.
- Nedeljkovic, M., Bouzas-Ammann, G., Zimmermann, L., Stute, P., Ausfeld-Hafter, B., 2013. Modalities of acupuncture treatments in assisted reproductive technology – a comparison of treatment practice in Swiss, German, and Austrian fertility centers with findings from randomized controlled trials. *Forsch. Komplementmed.* 20, 112–118.
- Noll, A.A., Wilm, S., 2009. *Chinese Medicine in Fertility Disorders*. Thieme, Stuttgart, Germany.
- Olivennes, F., Alvarez, S., Bouchard, P., Fanchin, R., Salat-Baroux, J., Frydman, R., 1998. The use of a GnRH antagonist (Cetrorelix) in a single dose protocol in IVF-embryo transfer: a dose finding study of 3 versus 2 mg. *Hum. Reprod.* 13, 2411–2414.
- Paulus, W.E., Zhang, M., Strehler, E., El-Danasouri, I., Sterzik, K., 2002. Influence of acupuncture on the pregnancy rate in patients who undergo assisted reproduction therapy. *Fertil. Steril.* 77, 721–724.
- Rashidi, B.H., Tehrani, E.S., Hamedani, N.A., Pirzadeh, L., 2013. Effects of acupuncture on the outcome of in vitro fertilisation and intracytoplasmic sperm injection in women with polycystic ovarian syndrome. *Acupunct. Med.* 31, 151–156.
- Ried, K., Stuart, K., 2011. Efficacy of traditional Chinese herbal medicine in the management of female infertility: a systematic review. *Complement. Ther. Med.* 19, 319–331.
- Ritenbaugh, C., Verhoef, M., Fleishmann, S., Boon, H., 2003. Whole Systems Research: a discipline for studying complementary and alternative medicine. *Altern. Ther. Health. Med.* 9, 32–36.
- SART and ASRM, 2006. Guidelines on number of embryos transferred. *Fertil. Steril.* 86, S51–S52.
- SART and ASRM, 2008. Guidelines on number of embryos transferred. *Fertil. Steril.* 90, S163–S164.
- SART and ASRM, 2009. Guidelines on number of embryos transferred. *Fertil. Steril.* 92, 1518–1519.
- Shen, C., Wu, M., Shu, D., Zhao, X., Gao, Y., 2015. The role of acupuncture in in vitro fertilization: a systematic review and meta-analysis. *Gynecol. Obstet. Invest.* 79, 1–12.
- Smith, C., Coyle, M., Norman, R.J., 2006. Influence of acupuncture stimulation on pregnancy rates for women undergoing embryo transfer. *Fertil. Steril.* 85, 1352–1358.
- Smith, C.A., Zhu, X., He, L., Song, J., 2011. Acupuncture for primary dysmenorrhoea. *Cochrane Database Syst. Rev.* (1), CD007854.
- Smith, J.F., Eisenberg, M.L., Millstein, S.G., Nachtigall, R.D., Shindel, A.W., Wing, H., Cedars, M., Pasch, L., Katz, P.P., 2010. The use of complementary and alternative fertility treatment in couples seeking fertility care: data from a prospective cohort in the United States. *Fertil. Steril.* 93, 2169–2174.

- StataCorp, 2011. Stata Statistical Software: Release 12. College Station, TX.
- Stener-Victorin, E., 2005. The pain-relieving effect of electro-acupuncture and conventional medical analgesic methods during oocyte retrieval: a systematic review of randomized controlled trials. *Hum. Reprod.* 20, 339–349.
- Stener-Victorin, E., Waldenström, U., Andersson, S.A., Wikland, M., 1996. Reduction of blood flow impedance in the uterine arteries of infertile women with electro-acupuncture. *Hum. Reprod.* 11, 1314–1317.
- Stener-Victorin, E., Baghaei, F., Holm, G., Janson, P.O., Olivecrona, G., Lönn, M., Mannerås-Holm, L., 2012. Effects of acupuncture and exercise on insulin sensitivity, adipose tissue characteristics, and markers of coagulation and fibrinolysis in women with polycystic ovary syndrome: secondary analyses of a randomized controlled trial. *Fertil. Steril.* 97, 501–508.
- Tan, L., Tong, Y., Sze, S.C., Xu, M., Shi, Y., Song, X.Y., Zhang, T.T., 2012. Chinese herbal medicine for infertility with anovulation: a systematic review. *J. Altern. Complement. Med.* 18, 1087–1100.
- Vernon, M., Stern, J.E., Ball, G.D., Wininger, D., Mayer, J., Racowsky, C., 2011. Utility of the national embryo morphology data collection by the Society for Assisted Reproductive Technologies (SART): correlation between day-3 morphology grade and live-birth outcome. *Fertil. Steril.* 95, 2761–2763.
- Vickers, A.J., Cronin, A.M., Maschino, A.C., Lewith, G., Macpherson, H., Foster, N.E., Sherman, K.J., Witt, C.M., Linde, K., 2012. Acupuncture for chronic pain: individual patient data meta-analysis. *Arch. Intern. Med.* 172, 1444–1453.
- Westergaard, L.G., Mao, Q., Krogslund, M., Sandrini, S., Lenz, S., Grinsted, J., 2006. Acupuncture on the day of embryo transfer significantly improves the reproductive outcome in infertile women: a prospective, randomized trial. *Fertil. Steril.* 85, 1341–1346.
- WHO, 2007. Western Pacific Region WHO international Standard Terminologies on Traditional Medicine in the Western Pacific Region. World Health Organization, Geneva, Switzerland.

Declaration: Support provided by NIH/NCCAM grant R25 AT002879 (Suppl) and 1K23AT006392. LHR owned the acupuncture practice. The authors report no financial or commercial conflicts of interest.

Received 11 November 2014; refereed 6 February 2015; accepted 10 February 2015.